



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Corrigendum

Citation for published version:

Alfituri, OA, Quintana, JF, MacLeod, A, Garside, P, Benson, RA, Brewer, JM, Mabbott, NA, Morrison, LJ & Capewell, P 2021, 'Corrigendum: To the Skin and Beyond: The Immune Response to African Trypanosomes as They Enter and Exit the Vertebrate Host', *Frontiers in Immunology*, vol. 12, 780758. <https://doi.org/10.3389/fimmu.2021.780758>

Digital Object Identifier (DOI):

[10.3389/fimmu.2021.780758](https://doi.org/10.3389/fimmu.2021.780758)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Frontiers in Immunology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Corrigendum: To the Skin and Beyond: The Immune Response to African Trypanosomes as They Enter and Exit the Vertebrate Host

Omar A. Alfituri¹, Juan F. Quintana², Annette MacLeod², Paul Garside², Robert A. Benson², James M. Brewer², Neil A. Mabbott¹, Liam J. Morrison¹ and Paul Capewell^{3*}

¹ Roslin Institute, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, United Kingdom, ² Wellcome Centre for Integrative Parasitology, Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom, ³ College of Medical, Veterinary and Life Sciences, Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow, United Kingdom

OPEN ACCESS

Edited and reviewed by:

Yasuyuki Goto,
The University of Tokyo, Japan

*Correspondence:

Paul Capewell
paul.capewell@glasgow.ac.uk

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Immunology

Received: 21 September 2021

Accepted: 11 October 2021

Published: 27 October 2021

Citation:

Alfituri OA, Quintana JF, MacLeod A, Garside P, Benson RA, Brewer JM, Mabbott NA, Morrison LJ and Capewell P (2021) Corrigendum: To the Skin and Beyond: The Immune Response to African Trypanosomes as They Enter and Exit the Vertebrate Host. *Front. Immunol.* 12:780758. doi: 10.3389/fimmu.2021.780758

Keywords: African trypanosomiasis, *Trypanosoma brucei*, skin, transmission, innate immunity, neglected tropical disease

A Corrigendum on:

To the Skin and Beyond: The Immune Response to African Trypanosomes as They Enter and Exit the Vertebrate Host

by Alfituri OA, Quintana JF, MacLeod A, Garside P, Benson RA, Brewer JM, Mabbott NA, Morrison LJ and Capewell P (2020). *Front. Immunol.* 11:1250. doi: 10.3389/fimmu.2020.01250

In the original article, there was a mistake in **Figure 1** as published. Panel C had an incorrectly orientated lymph node schema. The corrected **Figure 1** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Alfituri, Quintana, MacLeod, Garside, Benson, Brewer, Mabbott, Morrison and Capewell. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 1 | The skin, draining lymphatics, and lymph nodes. **(A)** Diagram of the cellular composition of the epidermal, dermal, and subcutaneous layers of mammalian skin. The outermost epidermal layer consists of a layer of keratinocytes above a layer of keratinocytes. These cells manage the tight junctions and the stratum corneum. Langerhans cells and intraepithelial T cells survey the epidermis for antigen to be presented. The central dermal layer contains fibroblasts that produce extracellular matrix proteins to provide structural support and elasticity. Immune responses are initiated by dermal macrophages, dermal dendritic cells, NK cells, and T cells. The inner subcutaneous layer primarily consists of adipocytes. Local lymphatic and blood vessels allow for the trafficking of cells, proteins, and waste. The initial tsetse fly bite injects trypanosomes into the dermis. From the dermis, the parasites exhibit tropism that leads to migration toward the afferent lymph vessels in the skin disseminating to the blood and other regions of the body. **(B)** The mechanism behind directional migration of trypanosomes from the skin to the lymphatics is unknown. Parasites may be responding to an unreported chemical cue in a chemotactic manner and they may crawl along lymph vessels, access open junctions, or are drawn into the lymphatics through hydrodynamic flow force and pressure. **(C)** Afferent lymphatic vessels in the skin allow for the drainage of leukocytes and antigen into the draining lymph node. Lymph, containing activated T and B cells, plasma cells, and antibody, passes into the medullary sinus, before exiting via efferent lymphatic vessels. Trypanosomes enter the draining lymph nodes, causing lymphadenopathy, and exit via the efferent lymphatics. Systemic dissemination of the host is reached via the main lymphatic ducts. The figure uses art adapted from material provided by Servier Medical Art under a Creative Commons Attribution 3.0 Unported License (<https://creativecommons.org/licenses/by/3.0/>).